

The Lattice Dimension of a Tree

Sergei Ovchinnikov
 Mathematics Department
 San Francisco State University
 San Francisco, CA 94132
 sergei@sfsu.edu

Abstract

The lattice dimension of a graph G is the minimal dimension of a cubic lattice in which G can be isometrically embedded. In this note we prove that the lattice dimension of a tree with n leaves is $\lceil n/2 \rceil$.

In the paper, T is a tree with q edges and n leaves. By the Djoković theorem [3] (see also [1, 2, 5]), T can be isometrically embedded into the q -cube $\mathcal{Q}_q$ and cannot be embedded into a cube of a smaller dimension. The number q is the *isometric dimension* $\dim_I(T)$ of the tree T [2]. Thus the vertices of T can be labeled with 0/1 q -dimensional vectors in such a way that the distance between two edges is the Hamming distance between the corresponding vectors. This labeling is also known as an *addressing scheme* for T [1].

Let $\mathbb{Z}^d$ be the edge skeleton of the standard cubic lattice of dimension d . Simple examples show that, generally speaking, T can be isometrically embedded into $\mathbb{Z}^d$ of a smaller dimension than $\dim_I(T)$. In a more general setting, one can consider the following problem: for a given graph G , find the minimum possible dimension d such that G can be isometrically embedded into $\mathbb{Z}^d$. We call this number d the *lattice dimension* of G and denote it by $\dim_Z(G)$. Recently, Eppstein [4] showed how to express the lattice dimension of a partial cube G in terms of a maximal matching in the semicube graph $\text{Sc}(G)$ associated with G , and presented a polynomial time algorithm for finding this dimension. In this paper, we prove directly that the lattice dimension of a tree with n leaves is $\lceil n/2 \rceil$.

Clearly, a tree with two leaves is embeddable into $\mathbb{Z}$. In what follows T is a tree with $n > 2$ leaves. For a given leaf $v \in T$ we denote v_T the closest vertex in T which has a degree greater than two and call the unique path (v_T, v) the *hanging path* to v .

Lemma 1. *Suppose the star $K_{1,n}$ is isometrically embedded into $\mathbb{Z}^d$. Then*

$$d \geq \lceil n/2 \rceil,$$

Moreover, $\dim_Z(K_{1,n}) = \lceil n/2 \rceil$.

Proof. We may assume that the center of $K_{1,n}$ is represented by the zero vertex in $\mathbb{Z}^d$. Then each leaf in $K_{1,n}$ is represented by the end vertex of a coordinate unit vector e_k or $-e_k$. It follows that $2d \geq n$. Since d is an integer, we have $d \geq \lceil n/2 \rceil$.

We construct an isometric embedding of $K_{1,n}$ into $\mathbb{Z}^{\lceil n/2 \rceil}$ as follows. Let $v_1, \dots, v_n$ be the leaves and c be the center of $K_{1,n}$. We map c into the zero vertex of $\mathbb{Z}^{\lceil n/2 \rceil}$, v_1 and v_2 into the end points of e_1 and $-e_1$, respectively, and so on. Clearly, we obtain an isometric embedding of $K_{1,n}$ into $\mathbb{Z}^{\lceil n/2 \rceil}$. $\square$

Lemma 2. *Let T be a tree with n leaves. Suppose that T is isometrically embedded into $\mathbb{Z}^d$. Then*

$$d \geq \lceil n/2 \rceil.$$

Proof. Let (u, v) be an inner edge of T , that is an edge with vertices that are not leaves. Vertices u and v are represented by vertices in $\mathbb{Z}^d$ that are different only in one position, say,

$$u = (x_1, x_2, \dots, x_d), \quad v = (x'_1, x_2, \dots, x_d),$$

with $|x_1 - x'_1| = 1$. Let us select all edges in $\mathbb{Z}^d$ with vertices that have the same first coordinates as u and v . Since T is a tree and the embedding is isometric, the edge (u, v) is the only edge among selected which belongs to T . Now we remove all the selected edges in $\mathbb{Z}^d$ and, for each selected edge, identify vertices defining this edge. We obtained a tree which is a contraction [5] of T having the same number n of leaves and embedded into another copy of $\mathbb{Z}^d$. By repeating this process, we end up with the star $K_{1,n}$ embedded into $\mathbb{Z}^d$. By Lemma 1, we have $d \geq \lceil n/2 \rceil$. $\square$

Now we prove the main theorem of the paper.

Theorem 1. *The lattice dimension of a tree T with n leaves is $\lceil n/2 \rceil$.*

Proof. By Lemma 2, it suffices to show that there is an isometric embedding of T into $\mathbb{Z}^{\lceil n/2 \rceil}$. To construct this embedding we use an inductive argument.

Suppose that the statement of the theorem is true for all trees with the number of leaves less than n for a given $n > 2$. Let v and u be two leaves of T and (v_T, v) and (u_T, u) be their hanging paths. The degrees of vertices v_T and u_T are greater than two and these two vertices are not necessarily different.

Suppose first that $v_T \neq u_T$. By removing the hanging paths (u_T, u) and (v_T, v) , we obtain a new tree T' with $(n-2)$ leaves. By the induction hypothesis, T' can be isometrically embedded into $\mathbb{Z}^d$ with $d = \lceil \frac{n-2}{2} \rceil = \lceil \frac{n}{2} \rceil - 1$. Now we embed, in a natural way, $\mathbb{Z}^d$ into $\mathbb{Z}^{d+1}$ and add the paths (u_T, u) and (v_T, v) to the image of T' in the positive and negative directions of the new dimension, respectively. Clearly, we constructed an isometric embedding of T into $\mathbb{Z}^{\lceil n/2 \rceil}$.

Suppose now that all leaves v have the same vertex $c = v_T$ in paths (v_T, v) . Then c is the only vertex in T of degree greater than two. By modifying in an obvious way the construction from Lemma 1, we obtain the required embedding. $\square$

Suppose T is isometrically embedded into $\mathbb{Z}^d$ with $d = \lceil n/2 \rceil$. The projection of T into the k 'th coordinate axis is a path P_{l_k} of length $l_k > 0$. Thus T is actually embedded into the *complete grid graph* [5]

$$P = P_{l_1} \square P_{l_2} \square \cdots \square P_{l_d},$$

where $\square$ denotes the Cartesian product. We have

$$d = \dim_Z(T) = \dim_Z(P) \quad \text{and} \quad q = \dim_I(T) = \dim_I(P).$$

Although the grid P is not uniquely defined by T , the set $\mathcal{L} = \{l_1, \dots, l_d\}$ determines both the lattice and isometric dimensions of T . Namely,

$$\dim_Z(T) = |\mathcal{L}| \quad \text{and} \quad \dim_I(T) = l_1 + \cdots + l_d.$$

Clearly, we may assume that $P_{l_k} = \{0, 1, \dots, l_k\}$. Then an isometric embedding of T into P defines an addressing scheme for T in which each vertex is labeled by a d -dimensional vector with the k 's coordinate ranging from 0 to l_k . The distance between two edges is the ℓ_1 -distance between the corresponding vectors.

References

- [1] A.S. Asartian, T.M.J. Denley, and R. Häggkvist, *Bipartite Graphs and their Applications* (Cambridge University Press, 1998).
- [2] M.M. Deza and M. Laurent, *Geometry of Cuts and Metrics* (Springer, 1997).
- [3] D.Z. Djoković, Distance preserving subgraphs of hypercubes, *J. Combin. Th. Ser. B* **14** (1973) 263–267.
- [4] D. Eppstein, The lattice dimension of a graph, Manuscript, 2004.
- [5] W. Imrich and S. Klavžar, *Product Graphs*, (John Wiley & Sons, 2000).